

DESERT LOCUST CONTROL ORGANIZATION FOR EASTERN AFRICA

..... DLCO-EA)



Headquarters (Addis Ababa)

Tel: 251-1-16461477/0287/0290

Fax: 251-1-16460296

Operations Office (Nairobi)


Tel: 254-020-6002305/6001488

Fax: 254-020-6001575

SITREP No. 10/2014-2015

DESERT LOCUST AND OTHER MIGRATORY PESTS SITUATION REPORT FOR

APRIL, 2015



1.0 WEATHER AND ECOLOGICAL CONDITIONS

In the Central Region, no significant rains fell during April. Consequently, ecological conditions continued to dry out in coastal areas along both sides of the Red Sea. Breeding conditions improved in the spring breeding areas of the interior of Saudi Arabia from good rains in late March. Light rains fell over parts of the northern Somali plateau at times during the second half of April. (*FAO DL bulletin No.439*)

1.1 Djibouti

Report not received.

1.2 Eritrea

Light rains fell during April mainly in the southern highlands of the country. Vegetation on the coastal areas has dried out during the month except for some which remained green in earlier flooded areas.

1.3 Ethiopia

Widespread rainfall has occurred in the southern and southwestern parts of the country during the first two weeks of April. By the end of the month, rainfall coverage has increased and the northern and some locations in the east have received low to moderate amount of rainfall.

Consequently, annual vegetation started greening in areas where rains fell.

1.4 Kenya

During April, moderate to heavy rains fell covering most parts of the country. Consequently, newspapers reported that many families have been displaced and infrastructure have been damaged by floods following some heavy rains that fell during the 3rd decade of the month in the western and the Rift Valley regions of the country. Annual and perennial vegetation remained green across most parts of the country.

1.5 Somalia

Localized low to moderate rains fell in the northwestern escarpments and the northern coastal areas during the second decade of April

Low to moderate rains also fell in the northeastern, the central and the southern parts of the country. Consequently, floods and river breakages were occurred in the middle Shabelle region.

Vegetation in most parts of the country was reported dry except for some, which remained green in flooded and irrigated areas.

Rainfall record (mm) during April, 2015

Date	Burao	Xudun	Galkayo	Garowe	Aburiin
14	15.00	0.00	0.00	0.00	0.00
15	0.00	9.00	0.60	0.00	0.00
16	0.00	25.00	0.00	0.00	0.00
17	0.00	0.00	9.20	0.00	0.00
22	0.00	0.00	21.00	0.00	0.00
24	0.00	0.00	0.00	0.00	3.80
25	5.60	0.00	0.00	0.00	0.00
26	1.40	0.00	55.40	14.60	0.00
Total	22.00	34.00	86.20	14.60	3.80

1.6 Sudan

No significant rains fell and vegetation dried out in most of the winter Desert Locust breeding areas along the Red Sea coast.

1.7 Tanzania

Medium to heavy rains fell during April covering most parts of the country. Consequently, annual and perennial vegetation were green in wider areas of the country.

1.8 Uganda

Moderate to heavy amount of rains fell during April. Consequently vegetation turned green across most parts of the country.

2.0 Desert Locust (*Schistocercagregaria*)

2.1 Djibouti

No locusts were reported.

2.2 Eritrea

Control operation that was going on since November, 2014 on the Red Sea coastal areas came to an end during April. However, some isolated locusts remained in the northern Red Sea coast mainly in irrigated farms.

2.3 Ethiopia

No locusts were reported.

2.4 Somalia

No locusts were reported.

2.5 Sudan

Red Sea State:

Low densities of immature and mature solitarious adults remained in the central and southern Red Sea coast mainly between Suakin and Tokar. Scattered mature and immature solitarious adults were also seen in cropping areas along the Atbara River. A DLCO-EA aircraft continued to be positioned in the country but no control operation was conducted.

Situation in Other Regions and Forecast

(Extracted from FAO DL Bulletin No. 439)

Central Region: low numbers of solitarious adults remained in a few places along the coast in Sudan and Saudi Arabia and no further control operations were required. A similar situation is likely on the northern and central Red Sea coast of Eritrea. Scattered adults were seen along the Atbara River in the interior of northern Sudan that probably arrived from the Red Sea coastal winter breeding areas. No locusts were reported elsewhere in the region. During the forecast period, small-scale breeding may occur in northern Sudan and in the interior of Saudi Arabia. the situation improved in the winter breeding areas along both sides of the Red Sea due to control operations and drying conditions in March. In Sudan, ground and aerial control operations declined, treating mainly locally bred adult groups and swarms on the southern coast. A few adult groups and swarms moved into this area from Eritrea where control was in progress against similar infestations. Locust numbers declined on the Red Sea coast in Saudi Arabia where limited control operations were conducted in the north. Low numbers of adults persisted on the Red Sea and Gulf of Aden coasts in Yemen.

Western Region: The situation remained calm in April. Only a few isolated solitarious locusts were present north of the Hoggar Mountains in the central Sahara of Algeria.

Eastern Region: The situation remained calm during April. Scattered solitarious adults were observed in a few places on the coast and in the interior of southeast Iran.

3.0 Forecast until mid-June, 2015

3.1 Djibouti

No significant developments are likely.

3.2 Eritrea

Low to moderate numbers of adults may be present in parts of the highlands.

3.3 Ethiopia

No significant developments are likely.

3.4 Somalia

No significant developments are likely.

3.5 Sudan

Scattered adults are likely to persist in and near cropping areas along the Nile and the Atbara Rivers in River Nile and Northern States where small-scale breeding may occur.

3.6 Kenya, Tanzania and Uganda

The countries are expected to remain free of Desert Locust infestations.

4.0 OTHER MIGRATORY PESTS

4.1 Red-billed Quelea birds (*Quelea quelea* sp.)

4.1.1 Kenya

Quelea birds outbreak was reported in Kisumu County where they were attacking irrigated Rice

4.1.2 Tanzania

Quelea birds outbreaks were reported in the country and a DLCO-EA aircraft has been deployed to control the infestations. Details of the infestation and

operations have not been received during the reporting period.

4.1.2 Ethiopia

No infestation reported.

4.1.3 Eritrea

Report not received.

4.2 African Armyworm (*Spodoptera exempta*)

No infestation reported in the region during April.

Forecast for May, 2015

It is likely that early infestation and outbreak to occur mainly in the southern and southeastern parts of Ethiopia that could extend to the mid-Rift Valley areas. Therefore, it is highly recommended that monitoring of moths and assessment of early outbreaks to continue in the suspected traditional breeding and migration locations.

4.3 Tsetse fly

4.3.1 Uganda

Infestation not reported.

CIFO

For Director,

7th May, 2015

For more information about the Organization, please visit DLCO-EA's Website:

www.dlcoea.org.et